

First record of the White-winged Vampire bat, *Diaemus youngii* (Jentink, 1893) (Chiroptera, Phyllostomidae) for the state of Goiás, Brazil, with a revised distribution map

João Paulo Maires Hoppe¹, Gabriella Marchezi², Albert David Ditchfield²

1 Laboratório de Biodiversidade de Insetos (LaBI), Departamento de Ciências Biológicas, Universidade Federal do Espírito Santo (UFES), Avenida Fernando Ferrari, 514, Edifício Lydia Behar, Bloco A, Sala 213, Vitória, Espírito Santo, CEP 29075.970, Brazil. **2** Laboratório de Estudos em Quirópteros (LABEQ), Departamento de Ciências Biológicas, Universidade Federal do Espírito Santo (UFES), Avenida Fernando Ferrari, 514, Edifício Lydia Behar, Bloco A, Sala 309, Vitória, Espírito Santo, CEP 29075.970, Brazil.

Corresponding author: João Paulo Maires Hoppe, jpmhoppe@gmail.com

Abstract

Diaemus youngii (Jentink, 1893) is a widespread species occurring from southern Mexico to northern Argentina. Along its distribution, however, this bat is rare to uncommon. Here, we report its record for the state of Goiás, central-western Brazil, based on a specimen found in the mammal collection of the Instituto Nacional da Mata Atlântica, former Museu de Biologia Professor Mello Leitão. *Diaemus youngii* is known from 129 unique localities, with 81 (over 62%) in Brazil, followed by Peru (7), Bolivia (6) and Venezuela (6). It is absent in the west of the Andes, southern Argentina, Uruguay, and Caribbean islands, except for Trinidad.

Key words

Cerrado, Desmodontinae, Neotropical region, scientific collection.

Academic editor: Marcelo Nogueira | Received 7 September 2018 | Accepted 19 December 2018 | Published 18 January 2019

Citation: Hoppe JPM, Marchezi G, Ditchfield AD (2019) First record of the White-winged Vampire bat, *Diaemus youngii* (Jentink, 1893) (Chiroptera, Phyllostomidae) for the state of Goiás, Brazil, with a revised distribution map. Check List 15 (1): 55–64. <https://doi.org/10.15560/15.1.55>

Introduction

Bats are still widely held as evil, disease-carrying, blood-sucking animals, especially in the western world, where old myths persist (Teixeira and Papavero 2003). However, only 3 among the more than 1,300 extant bat species feed on blood: *Desmodus rotundus* (É. Geoffroy Saint-Hilaire, 1810), *Diphylla ecaudata* Spix, 1823, and *Diaemus youngii* (Jentink, 1893). These 3 species belong to the Desmodontinae subfamily and are endemic of the Americas. These bats also share several adaptations that allow the consumption and digestion of blood: sharp, canine-like incisors, saliva with anticoagulant

components, and a sac-like elastic stomach, specialized for liquid absorption (Kwon and Gardner 2008). Due to their feeding habits, vampire bats are of epidemiological concern, especially for their role in the transmission of rabies in rural areas, affecting humans and other mammals (Calisher et al. 2006, Dantas-Torres 2008).

Diaemus youngii has a wide distribution, occurring from Mexico to northern Argentina (Kwon and Gardner 2008). Along its distribution, however, it is a rare to uncommon species (Aguiar et al. 2006). In Brazil, few records are available for most states where this bat has been found, and some well-studied states still lack records (e.g., Espírito Santo, Mendes et al. 2010). While




Figure 1. Known localities of *Diaemus youngii*. The star points the new occurrence at Monte Alegre de Goiás, state of Goiás, Brazil. See supplementary material for coordinates and literature sources.

examining material in a museum mammal collection, we discovered a specimen of *D. youngii* that fills the gap for the state of Goiás. Here we report this finding and provide an updated map covering the whole distribution of the species.

Methods

The specimen of *D. youngii* is housed in the mammal collection of the Instituto Nacional da Mata Atlântica, Santa Teresa (INMA), Brazil, formerly known as Museu de Biologia Professor Mello Leitão. Craniodental measurements were obtained using a digital caliper calibrated to the nearest 0.01 mm, following the protocol devised by Vizotto and Taddei (1973). We gathered all available published records of *D. youngii* from the literature, giving preference to the primary sources (see Appendix).

Results

New records and distribution map. *Diaemus youngii*, MBML 2005, previously ISDF 1047, adult female,

orchard next to a poultry farm, Monte Alegre de Goiás, Goiás, Brazil (13°15'23" S, 046°54'02" W). The species is known from 129 unique localities from 16 countries, with Brazil (81 localities), Peru (7), Bolivia (6) and Venezuela (6) being the most represented (Fig. 1). Craniodental measurements (Table 1). A. Bredt, E. Magalhães, P.H. Oliveira, 25 April 1995.

Identification. We used the identification keys of Kwon and Gardner (2008) and Díaz et al. (2016) to identify our specimen. *Diaemus youngii* can be differentiated from *De. rotundus* and *Di. ecaudata*, by presenting moderately elongated thumbs with 1 basal pad, white wing-tips, and 2 upper molars on each side (Greenhall and Schutt 1996). Our specimen was confidently identified based on these characteristics (Fig. 2), and overall agrees with descriptions in the literature. The inner lower incisors are described as trilobated (Kwon and Gardner 2008), but in our specimen these teeth are worn down (Fig. 2A). Notwithstanding, Greenhall and Schutt (1996) have noticed that the lower incisors in *Diaemus* tend to vary in kind and degree of lobation. Mensural data from our specimen are close to those reported for the holotype (Carter and Dolan 1978), and for specimens from

central-western and northern Brazil (Pedroso et al. 2018), and Venezuela and Trinidad (Greenhall and Schutt 1996) (Table 1). These data suggest little size variation within *D. youngii*, but larger samples are required to further investigate this aspect.

Discussion

Desmodontinae are known to feed on mammalian and avian blood, with *Di. ecaudata* regarded as an avian blood specialist, while *De. rotundus* and *D. youngii* feed on a broader spectrum (Ito et al. 2016). However, *D. youngii* seems to prefer avian blood, perhaps to avoid competition of resources with *De. rotundus* (Gardner 1977, Sazima and Uieda 1980, Greenhall and Schutt 1996). Our specimen was collected in an orchard, but next to a poultry farm containing a few dozen chickens. Unfortunately, the records did not mention if the fowl presented bite marks or signs of decaying health.

Vampire bats are habitually exposed to blood, even by sharing blood meals with conspecifics in their roost places (Wilkinson 1984), increasing the odds of acquiring rabies virus (RABV). RABV are frequently isolated from *De. rotundus*, but less commonly in other hematophagous bats (Castilho et al. 2010). There are a few positive RABV isolates from *D. youngii*, with reports from the Brazilian states of Piauí and Maranhão (Castilho et al. 2010, Póvoas et al. 2012), and also Mexico and Trinidad and Tobago (Escobar et al. 2015). Before donation to INMA, the specimen reported here was screened for RABV by the Instituto de Saúde do Distrito Federal (ISDF, number 1047), with negative results. *Desmodus rotundus* can efficiently digest mammal blood and was thought to be the only vampire bat preying on humans. However, there are at least 2 records of *Di. ecaudata*, an avian blood specialist with physiological adaptations similar to *D. youngii* (see Ito et al. 2016), feeding on humans (Ruschi 1953, Ito et al. 2016).

Housing conditions may influence the foraging behavior of vampire bats, especially in rural or poor areas where domestic animals, often maintained together, are in close proximity with human facilities. *Diaemus youngii* has also been captured in urban areas (Urbieta et al. 2017). These conditions favor the transmission of RABV, even if indirectly (Turner 1975).

According to the literature, *D. youngii* has a wide distribution in the Neotropical region, occurring from southern Mexico to northern Argentina. It seems to be absent, however, from the west of the Andes and most Caribbean islands (except for Trinidad). Suriname, Chile, and Uruguay, in South America, and Belize and Honduras, in Central America also lack records (Fig. 1; see also Greenhall and Schutt 1996). The distribution of this species seems to be influenced by thermoregulation issues (e.g. intolerance to colder climates), shared with *Diphylla*, but not with *Desmodus* (McNab 1969). The latter genus of bat has a greater ability to maintain its body temperature, and its distribution is known to reach as far as Chile and Uruguay (Acosta y Lara 1950, Delpietro et al. 2017). The absence of *D. youngii* in Honduras (Goodwin 1942, Valdez and LaVal 1971), Belize (Fenton et al. 2001), and Suriname, however, may represent sampling gaps. In northern South America, all countries bordering Suriname have reported its occurrence (Piccinini 1974, Peracchi et al. 1984, Bernard and Fenton 2002).

The large number of records of *D. youngii* in southeastern Brazil, especially in Rio de Janeiro and São Paulo, can be attributed to the long-duration surveys conducted in those states (Esbérard and Bergallo 2005, Garbino 2016). Contrasting with the situation in this region, however, the majority of the Brazilian territory is poorly sampled (Bernard et al. 2011), including the northeastern region. Some efforts have been made recently in this region, but no record of *D. youngii* is currently available from Rio Grande do Norte (Vargas-Mena et al. 2018), Ceará (Silva et al. 2015) and Roraima (Capaverde Junior

Table 1. Craniodental measurements (in mm) of *Diaemus youngii* from Monte Alegre de Goiás, Goiás, Brazil (MBML 2005), and comparative data for the holotype (Upper Canje Creek, Guyana; RNH 12088; Carter and Dolan 1978), and for material from central-western and northern Brazil (Pedroso et al. 2018), and from Trinidad and Venezuela (Greenhall and Schutt 1996).

| Variables | MBML 2005 (GO) | Holotype | Greenhall and Schutt 1996 [†] | Pedroso et al. 2018 [‡] |
|---------------------------|----------------|----------|--|----------------------------------|
| Greatest length of skull | 24.35 | 25.0 | 24.8 | 25.61 |
| Condylobasal length | 21.58 | 23.0 | — | 21.01 |
| Condyllo-canine length | 19.63 | — | — | — |
| Basal length | 19.63 | — | — | — |
| Palatal length | 8.72 | — | — | 7.84 |
| Upper toothrow length | 3.10 | 3.5 | 3.5 | 5.61 |
| Width across upper molars | 6.28 | — | — | 6.58 |
| Postorbital length | 6.34 | 6.5 | 6.2 | 6.32 |
| Braincase breadth | 13.17 | 13.5 | 13.0 | 13.02 |
| Zygomatic breadth | 13.91 | 14.3 | 13.9 | 13.99 |
| Mandibular length | 14.70 | 15.2 | — | 15.15 |
| Lower toothrow length | 4.17 | 4.2 | — | — |

[†]Mean values for 3 males.

[‡]Mean values for 5 females.




Figure 2. *Diaemus youngii* from Goiás state, Brazil (MBML 2005). **A.** Lower incisors. **B.** Right thumb, showing the single basal pad. **C.** Right wing, depicting the white wingtip characteristic of this species.

et al. 2014). Absences from the southern states of Rio Grande do Sul and Santa Catarina, on the other hand, may be related to their colder climates, as previously reported for Uruguay (Acosta y Lara 1950).

The new record of *D. youngii* reported here covers a gap in the middle of the Cerrado, with the closest localities known to harbor this species being, respectively,

Brazlândia, Distrito Federal (280 km north; Aguiar et al. 2006) and Estação Ecológica Serra Geral do Tocantins, Jalapão (300 km south; Gregorin et al. 2011).

The first director of the Museum of Vertebrate Zoology (University of California), Joseph Grinnell, wrote in 1910 on the importance of museum collections (Grinnell 1910):

It will be observed, then, that our efforts are not merely to accumulate as great a mass of animal remains as possible. On the contrary, we are expending even more time than would be required for the collection of the specimens alone, in rendering what we do obtain as permanently valuable as we know how, to the ecologist as well as to the systematist. It is quite probable that the facts of distribution, life history, and economic status may finally prove to be of more far-reaching value, than whatever information is obtainable exclusively from the specimens themselves.

Unfortunately, without proper care and maintenance, these fundamental repositories of biodiversity can be tragically lost, as recently seen in the Museu Nacional do Rio de Janeiro and in the Instituto Butantan, both in Brazil. It is widely recognized that most species in the world have yet to be described (Mora et al. 2011), and a significant part is already waiting for description in a museum collection (Fontaine et al. 2012). Our report, based on a specimen collected more than 20 years ago, reinforces the role of collections, in helping with the basic knowledge of “where” a species can be found.

Acknowledgements

We are grateful to Juliana Paulo da Silva at Instituto Nacional da Mata Atlântica for access to the collection and specimens. Marcelo Nogueira, Robert Forsyth, and an anonymous reviewer gave substantial contributions that greatly improved the manuscript. JPMH received a postgraduate scholarship from CAPES.

Authors' Contributions

JPMH examined and identified the specimen, prepared the map and figures. GM took cranial measurements. All authors wrote, revised, and approved the manuscript.

References

Acosta y Lara EF (1950) Quirópteros del Uruguay. Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 3: 1–71.

Aguiar LM de S, Camargo WR, Portella A de S (2006) Occurrence of white-winged vampire bat, *Diaemus youngi* (Mammalia, Chiroptera), in the Cerrado of Distrito Federal, Brazil. *Revista Brasileira de Zoologia* 23: 893–896. <http://doi.org/10.1590/s0101-81752006000300041>

Aguiar LM de S, Antonini Y (2016) Prevalence and intensity of Streblidae in bats from a Neotropical savanna region in Brazil. *Folia Parasitologica* 63: 1–8. <https://doi.org/10.14411/fp.2016.024>

Anderson S, Koopman KF, Creighton GK (1982) Bats of Bolivia: An annotated checklist. *American Museum Novitates* 2750: 1–24.

Barnett AA, Sampaio EM, Kalko EK V., Shapley RL, Fischer EA, Camargo G, Rodríguez-Herrera B (2006) Bats of Jaú National Park, central Amazônia, Brazil. *Acta Chiropterologica* 8: 103–128. [https://doi.org/10.3161/1733-5329\(2006\)8\[103:BOJNPC\]2.0.CO;2](https://doi.org/10.3161/1733-5329(2006)8[103:BOJNPC]2.0.CO;2)

Barquez RM (1984) Significativa extension del rango de distribucion de *Diaemus youngii* (Yentink, 1893) (Mammalia, Chiroptera, Phyllostomidae). *Historia Natural* 4: 67–68.

Barquez RM, Sánchez MS, Sandoval ML (2011) Nuevos registros de

murciélagos (Chiroptera) en el norte de Argentina. *Mastozoología Neotropical* 18: 11–24.

Barros JH da S, Romijn PC, Baptista C, Pinto AG de S, Madeira M de F (2008) Relato de infecção natural de morcegos por flagelados tripanosomatídeos em diferentes municípios do estado do Rio de Janeiro. *Revista da Sociedade Brasileira de Medicina Tropical* 41: 683–685. <https://doi.org/10.1590/S0037-86822008000600025>

Bernard E (2001) Vertical stratification of bat communities in primary forests of Central Amazon, Brazil. *Journal of Tropical Ecology* 17: 115–126. <http://doi.org/10.1017/S0266467401001079>

Bernard E, Aguiar LMS, Machado RB (2011) Discovering the Brazilian bat fauna: a task for two centuries? *Mammal Review* 41: 23–39. <http://doi.org/10.1111/j.1365-2907.2010.00164.x>

Bernard E, Fenton MB (2002) Species diversity of bats (Mammalia: Chiroptera) in forest fragments, primary forests, and savannas in central Amazonia, Brazil. *Canadian Journal of Zoology* 80: 1124–1180. <http://doi.org/10.1139/Z02-094>

Bobrowiec PED (2012) A Chiroptera preliminary survey in the middle Madeira River region of Central Amazonia, Brazil. *Mammalia* 76: 277–283. <https://doi.org/10.1515/mammalia-2011-0065>

Bolzan DP, Lourenço EC, Costa L de M, Luz JL, Jordão-Nogueira T, Dias D, Esbérard CEL, Peracchi AL (2010) Morcegos da região da Costa Verde e adjacências, litoral sul do estado do Rio de Janeiro. *Chiroptera Neotropical* 16: 585–594.

Bordignon MO, Shapiro JT (2018) Bat diversity in the western Brazilian Pantanal. *Mammalia* 82: 256–265. <https://doi.org/10.1515/mammalia-2016-0107>

Bowles JB, Cope JB, Cope EA (1979) Biological studies of selected Peruvian bats of Tingo María, Departamento de Huánuco. *Transactions of the Kansas Academy of Sciences* 82: 1–10.

Brennan JM (1967) New records of chiggers from the West Indies. *Studies on the Fauna of Curaçao and other Caribbean Islands* 24: 146–156.

Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T (2006) Bats: Important reservoir hosts of emerging viruses. *Clinical Microbiology Reviews* 19: 531–545. <http://doi.org/10.1128/cmr.00017-06>

Capaverde UD Jr, Pacheco SM, Duarte ME (2014) Murciélagos (Mammalia: Chiroptera) del área urbana del municipio de Boa Vista, Roraima, Brasil. *Barbastella* 7: 13–18.

Carter DC, Dolan PG (1978) Catalogue of type specimens of Neotropical bats in selected European museums. *Special Publications of the Museum Texas Tech University* 15: 1–136.

Castilho JG, Carnieli P, Oliveira R de N, Fahl W de O, Cavalcante R, Santana AA, Rosa WLGA, Carrieri ML, Kotait I (2010) A comparative study of rabies virus isolates from hematophagous bats in Brazil. *Journal of Wildlife Diseases* 46: 1335–1339. <http://doi.org/10.7589/0090-3558-46.4.1335>

Costa L de M, Oliveira DM, Dias e Fernandes AFP, Esbérard CEL (2008) Ocorrência de *Diaemus youngi* (Jentink 1893), Chiroptera, no estado do Rio de Janeiro. *Biota Neotropica* 8: 217–220.

Costa L de M, Esbérard CEL (2011) *Desmodus rotundus* (Mammalia: Chiroptera) on the southern coast of Rio de Janeiro state, Brazil. *Brazilian Journal of Biology* 71: 739–746.

Dantas-Torres F (2008) Bats and their role in human rabies epidemiology in the Americas. *Journal of Venomous Animals and Toxins including Tropical Diseases* 14: 193–202. <http://doi.org/10.1590/s1678-91992008000200002>

Delpietro HA, Russo RG, Carter GG, Lord RD, Delpietro GL (2017) Reproductive seasonality, sex ratio and philopatry in Argentina's common vampire bats. *Royal Society Open Science* 4: 160959. <http://doi.org/10.1098/rsos.160959>

Díaz MM, Solari S, Aguirre LF, Aguiar, LMS, Barquez RM (2016) Clave de Identificación de los Murciélagos de Sudamérica. *Publicación Especial N° 2, PCMA (Programa de Conservación de los Murciélagos de Argentina)*, 160 pp.

Esbérard CEL, Bergallo HG (2005) Research on bats in the state of Rio de Janeiro, southeastern Brazil. *Mastozoología Neotropical*

- 12: 237–243.
- Escobar LE, Peterson AT, Favi M, Yung V, Medina-Vogel G (2015) Bat-borne rabies in Latin America. *Revista do Instituto de Medicina Tropical de São Paulo* 57: 63–72. <http://doi.org/10.1590/s0036-46652015000100009>
- Falcão F de C (2007) Mammalia, Chiroptera, Phyllostomidae, *Diaemus youngi*: First record for the state of Bahia, northeastern Brazil. *Check List* 3: 330–332. <https://doi.org/10.15560/3.4.330>
- Feijó A, Araújo P, Fracasso MP de A, Santos KRP (2010) New records of three bat species for the Caatinga of the state of Paraíba, northeastern Brazil. *Chiroptera Neotropical* 16: 723–727.
- Feijó A, Nunes HL, Langguth A (2016) Mamíferos da Reserva Biológica Guaribas, Paraíba, Brasil. *Revista Nordestina de Biologia* 24: 57–74.
- Fenton MB, Bernard E, Bouchard S, Hollis L, Johnston DS, Lausen CL, Ratcliffe JM, Riskin DK, Taylor JR, Zigouris (2001) The bat fauna of Lamanai, Belize: roosts and trophic roles. *Journal of Tropical Ecology* 17: 511–524. <http://doi.org/10.1017/s0266467401001389>
- Fontaine B, Perrard A, Bouchet P (2012) 21 years of shelf life between discovery and description of new species. *Current Biology* 22: R943–R944. <http://doi.org/10.1016/j.cub.2012.10.029>
- Garbino GST (2016) Research on bats (Chiroptera) from the state of São Paulo, southeastern Brazil: annotated species list and bibliographic review. *Arquivos de Zoologia* 47: 43–128.
- Gardner AL (1977) Feeding habits. In: Baker RJ, Jones JK, Carter DC (Eds), *Biology of Bats of the New World Family Phyllostomidae*. Part II. Texas Tech Press, Lubbock, Texas, 293–350.
- Gardner AL, LaVal RK, Wilson DE (1970) The distributional status of some Costa Rican Bats. *Journal of Mammalogy* 51: 712–729. <https://doi.org/10.2307/1378297>
- Goodwin GG (1942) Mammals of Honduras. *Bulletin of the American Museum of Natural History* 79: 107–195.
- Gracioli G, Carvalho CJB (2001) Moscas ectoparasitas (Diptera, Hippoboscoidea) de morcegos (Mammalia, Chiroptera) do Estado do Paraná. II. Streblidae. Chave pictórica para gêneros e espécies. *Revista Brasileira de Zoologia* 18: 907–960. <https://doi.org/10.1590/S0101-81752001000300026>
- Greenbaum IF, Knox Jones J (1978) Noteworthy records of bats from El Salvador, Honduras, and Nicaragua. *Occasional Papers of the Museum Texas Tech University* 55: 1–7.
- Greenhall AM (1963) Use of mist nets and strychnine for vampire control in Trinidad. *Journal of Mammalogy* 44: 396–399. <https://doi.org/10.2307/1377208>
- Greenhall AM, Schutt WA (1996) *Diaemus youngi*. *Mammalian Species* 533: 1–7. <http://doi.org/10.2307/3504240>
- Gregorin R, Gonçalves E, Aires CC, Carmignotto AP (2011) Morcegos (Mammalia: Chiroptera) da Estação Ecológica Serra Geral do Tocantins: composição específica e considerações taxonômicas. *Biota Neotropica* 11: 299–311. <http://doi.org/10.1590/s1676-06032011000100028>
- Grinnell J (1910) The methods and uses of a research museum. *Popular Science Monthly* 77: 163–169.
- Handley Jr CO (1967) Bats of the Canopy of an Amazonian Forest. *Atas do Simpósio sobre a Biota Amazônica* 5: 211–215.
- Herrera HM, Rocha FL, Lisboa C V., Rademaker V, Mourão GM, Jansen AM (2011) Food web connections and the transmission cycles of *Trypanosoma cruzi* and *Trypanosoma evansi* (Kinetoplastida, Trypanosomatidae) in the Pantanal Region, Brazil. *Transactions of the Royal Society of Tropical Medicine and Hygiene* 105: 380–387. <https://doi.org/10.1016/j.trstmh.2011.04.008>
- Hutterer R, Verhaagh M, Diller J, Podlousky R (1995) An Inventory of Mammals Observed at Panguana Biological Station, Amazonian Peru. *Ecotropica* 1: 3–20.
- Ito F, Bernard E, Torres RA (2016) What is for dinner? First report of human blood in the diet of the hairy-legged vampire bat *Diphylla ecaudata*. *Acta Chiropterologica* 18: 509–515. <http://doi.org/10.3161/15081109acc2016.18.2.017>
- Jentink FA (1893) On a collection of bats from the West-Indies. *Notes from the Leyden Museum* 15: 278–283.
- Kalko EKV., Handley Jr CO (2001) Neotropical bats in the canopy: diversity, community structure, and implications for conservation. *Plant Ecology* 153: 319–333. <https://doi.org/10.1023/a:1017590007861>
- Kwon M, Gardner AL (2008) Subfamily Desmodontinae J. A. Wagner, 1840. In: Gardner AL (Ed.) *Mammals of South America. Marsupials, Xenarthrans, Shrews, and Bats*. Volume 1. The University of Chicago Press, 218–224.
- López-González C, Presley SJ, Owen RD, Willig MR, Fox IG (1998) Noteworthy records of bats (Chiroptera) from Paraguay. *Mastozoología Neotropical* 5: 41–45.
- Mares MA, Willig MR, Streilein KE, Lacher TE (1981) The mammals of northeastern Brazil: a preliminary assessment. *Annals of Carnegie Museum* 50: 81–137.
- Marques-Aguiar SA, Del Aguila M V., Aguiar GF de S, Saldanha N, Silva-Junior J de S, Rocha MMB (2003) Caracterização e perspectivas de estudo dos quirópteros da Estação Científica Ferreira Penna, município de Melgaço, Pará. In: Estação Científica Ferreira Penna - Dez Anos de Pesquisa na Amazônia, 1–3.
- Marques-Aguiar SA, Aguiar GF de S, Saldanha N, Rocha MMB, Fonseca RTDA, Reis-Filho VO (2009) Diversidade de morcegos (Mammalia: Chiroptera) no Baixo Rio Xingu, estado do Pará. In: *Anais do IX Congresso de Ecologia do Brasil*. São Lourenço-MG, 1–4.
- McBee K, Baker RJ, Pimm SL (1985) Utility of morphological distance measures and clustering algorithms: a test using phyllostomid bats. *Annals of Carnegie Museum* 54: 393–412.
- McNab BK (1969) The economics of temperature regulation in Neotropical bats. *Comparative Biochemistry and Physiology* 31: 227–268. [https://doi.org/10.1016/0010-406x\(60\)91651-x](https://doi.org/10.1016/0010-406x(60)91651-x)
- Menezes Jr LF, Duarte AC, Contilides MD, Peracchi AL (2015) Comparação da quiropteroфаuna em área florestada e área aberta da RPPN Fazenda Bom Retiro, Rio de Janeiro, Brasil. *Iheringia (Série Zoologia)* 105: 271–275. <https://doi.org/10.1590/1678-476620151053271275>
- Mendes P, Vieira TB, Oprea M, Lopes SR, Ditchfield AD, Zortéa M (2010) O conhecimento sobre morcegos (Chiroptera: Mammalia) do estado do Espírito Santo, sudeste do Brasil. *Papéis Avulsos de Zoologia* 50: 363–373.
- Miller GS (1906) Twelve new genera of bats. *Proceedings of the Biological Society of Washington* 19: 83–86.
- Mok WY, Lacey LA (1980) Algumas considerações ecológicas sobre morcegos vampiros na epidemiologia da raiva humana na Bacia Amazônica. *Acta Amazonica* 10: 335–342. <https://doi.org/10.1590/1809-43921980102335>
- Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B (2011) How many species are there on Earth and in the Ocean? *PLoS Biology* 9: e1001127. <http://doi.org/10.1371/journal.pbio.1001127>
- Nogueira MR, Pol A, Peracchi AL (1999) New records of bats from Brazil with a list of additional species for the chiropteran fauna of the state of Acre, western Amazon basin. *Mammalia* 63: 363–368. <https://doi.org/10.1515/mamm.1999.63.3.363>
- Olimpio APM, Ventura MC da S, Mascarenhas M de JO, Nascimento DC, Andrade FAG, Fraga E da C, Barros MC (2016) Bat fauna of the Cerrado savanna of eastern Maranhão, Brazil, with new species occurrences. *Biota Neotropica* 16: e20150089. <https://doi.org/10.1590/1676-0611-BN-2015-0089>
- Oliveira AKM, Oliveira MD, Favero S, Oliveira LF (2012) Diversity, similarity and trophic guild of chiropteroфаuna in three southern Pantanal sub-regions, state of Mato Grosso do Sul, Brazil. *Acta Scientiarum - Biological Sciences* 34: 33–39. <https://doi.org/10.4025/actascibiolsci.v34i1.7596>
- Osgood WH (1912) Mammals from western Venezuela and eastern Colombia. *Fieldiana: Zoology* 10: 33–66. <https://doi.org/10.5962/bhl.title.2647>
- Owen JG, Baker RJ, Jones JK (1990) First record of *Peromyscus*

- gymnotis* (Muridae) from El Salvador, with second records for *Choeroniscus godmani* and *Diaemus youngi* (Phyllostomidae). Texas Journal of Science 42: 417–418.
- Pacheco V, Cadenillas R, Velazco S, Salas E, Fajardo U (2007) Noteworthy bat records from the Pacific Tropical rainforest region and adjacent dry forest in northwestern Peru. Acta Chiropterologica 9: 409–422.
- Pedroso MA, Rocha PA, Brandão MV, Garbino GST, Moraes CO, Aires CC (2018) Filling gaps in the distribution of the white-winged vampire bat, *Diaemus youngii* (Phyllostomidae, Desmodontinae): New records for Southern Amazonia. Acta Amazonica 48: 154–157. <http://doi.org/10.1590/1809-4392201704291>
- Peracchi AL, Albuquerque ST (1971) Lista provisória dos quirópteros dos Estados do Rio de Janeiro e Guanabara, Brasil (Mammalia, Chiroptera). Revista Brasileira de Biologia 31: 405–413.
- Peracchi AL, Albuquerque ST (1986) Quirópteros do estado do Rio de Janeiro, Brasil (Mammalia, Chiroptera). Publicações Avulsas do Museu Nacional 66: 63–69.
- Peracchi AL, Raimundo SDL, Tannure AM (1984) Quirópteros do território federal do Amapá, Brasil (Mammalia, Chiroptera). Arquivos da Universidade Federal Rural do Rio de Janeiro 7: 89–100.
- Pereira SN, Dias D, Lima IP, Maas ACS, Martins MA, Bolzan DP, França DS, Oliveira MB, Peracchi AL, Ferreira MFS (2013) Mamíferos de um fragmento florestal em Volta Redonda, estado do Rio de Janeiro. Bioscience Journal 29: 1017–1027.
- Piccinini RS (1974) Lista provisória dos quirópteros da coleção do Museu Paraense Emílio Goeldi. Boletim do Museu Paraense Emílio Goeldi (Nova Série) 77: 1–32.
- Pinto AS, Bento DN (1986) *Trypanosoma cruzi*-like bloodstream trypanomastigotes in bats from the State of Piauí, Northeastern Brazil. Revista da Sociedade Brasileira de Medicina Tropical 19: 31–34. <https://doi.org/10.1590/S0037-86821986000100007>
- Pinto CM, Carrera JP, Mantilla-Meluk H, Baker RJ (2007) Mammalia, Chiroptera, Phyllostomidae, *Diaemus youngi*: First confirmed record for Ecuador and observations of its presence in museum collections. Check List 3: 244–247. <https://doi.org/10.15560/3.3.244>
- Póvoas DR, Chaves NP, Bezerra DC (2012) Raiva em quirópteros no estado do Maranhão: um estudo retrospectivo. Revista Brasileira de Ciência Veterinária 19: 163–166.
- Reis NR, Peracchi AL, Lima IP (2002) Morcegos da Bacia do Rio Tibagi. In: Medri ME, Bianchini E, Shibatta OA, Pimenta JA (Eds), A bacia do rio Tibagi. Londrina, 251–270.
- Reis NR, Barbieri ML da S, Lima IP, Peracchi AL (2003) O que é melhor para manter a riqueza de espécies de morcegos (Mammalia, Chiroptera): um fragmento florestal grande ou vários fragmentos de pequeno tamanho? Revista Brasileira de Zoologia 20: 225–230. <https://doi.org/10.1590/S0101-81752003000200009>
- Ruschi A (1953) Dois casos de sanguivorismo de *Desmodus rotundus* (E. Geoffroy) e *Diphylla ecaudata* Spix, no homem, e outras observações sobre os quirópteros hematófagos e acidentalmente hematófagos. Boletim do Museu de Biologia Mello Leitão (Biologia) 13: 1–8.
- Sanborn CC (1949) Mammals from the Rio Ucayali, Peru. Journal of Mammalogy 30: 277–288.
- Sá-Neto RJ, Marinho-Filho JS (2013) Bats in fragments of xeric woodland Caatinga in Brazilian semiarid. Journal of Arid Environments 90: 88–94. <https://doi.org/10.1016/j.jaridenv.2012.10.007>
- Sazima I, Uieda W (1980) Feeding behavior of the white-winged vampire bat, *Diaemus youngii*, on Poultry. Journal of Mammalogy 61: 102–104. <http://doi.org/10.2307/1379959>
- Schaller GB (1983) Mammals and their biomass on a Brazilian Ranch. Arquivos de Zoologia 31: 1–36.
- Sekiama ML, Reis NR, Peracchi AL, Rocha VJ (2001) Morcegos do Parque Nacional do Iguaçu, Paraná (Chiroptera, Mammalia). Revista Brasileira de Zoologia 18: 749–754. <https://doi.org/10.1590/S0101-81752001000300011>
- Silva C de L, Gracioli G (2013) Prevalence, mean intensity of infestation and host specificity of Spinturnicidae mites (Acari: Mesostigmata) on bats (Mammalia: Chiroptera) in the Pantanal, Brazil. Acta Parasitologica 58: 174–179. <https://doi.org/10.2478/s11686-013-0134-x>
- Silva LAM, Souza AQS, Lima AS, Araújo CSF, Silva CVM, Silva LEC, Gomes MF, Queiroz PL, Silva RM (2010) Morcegos da Reserva Particular do Patrimônio Natural Frei Caneca, nordeste do Brasil. In: V Encontro Brasileiro para o Estudo de Quirópteros. SBEQ, Búzios-RJ, 87–90.
- Silva SSP, Dias D, Martins MA, Guedes PG, Almeida JC, Cruz AP, Serra-Freire NM, Damascena JS, Peracchi AL (2015) Bats (Mammalia: Chiroptera) from the Caatinga scrublands of the Cratêus region, northeastern Brazil, with new records for the state of Ceará. Mastozoología Neotropical 22: 335–348.
- Sotero-Caio CG, Pieczarka JC, Nagamachi CY, Gomes AJB, Lira TC, O'Brien PCM, Ferguson-Smith MA, Souza MJ, Santos N (2011) Chromosomal homologies among vampire bats revealed by chromosome painting (Phyllostomidae, Chiroptera). Cytogenetic and Genome Research 132: 156–164. <https://doi.org/10.1159/000321574>
- Souza R de F, Novaes RLM, Siqueira AC, Sauwen C, Jacob G, Santos CEL, Felix S, Ribeiro E, Sant'Anna C, Vrcibradic D, Avilla L dos S, Sbragia IA, Santori RT (2015) Bats (Mammalia, Chiroptera) in a remnant of Atlantic Forest, Rio de Janeiro, southeastern Brazil. Neotropical Biology and Conservation 10: 9–14. <https://doi.org/10.4013/nbc.2015.101.02>
- Stutz WH, Albuquerque MC, Uieda W, Macedo EM, França CB (2004) Updated list of bats from Uberlândia, state of Minas Gerais, Southeastern Brazil. Chiroptera Neotropical 10: 188–190.
- Tavares V da C, Nobre CC, Palmuti CF de S, Nogueira E de PP, Gomes JD, Marcos MH, Silva RF, Farias SG, Bobrowiec PED (2017) The bat fauna from southwestern Brazil and its affinities with the fauna of western Amazon. Acta Chiropterologica 19: 93–106. <https://doi.org/10.3161/15081109ACC2017.19.1.007>
- Teixeira DM, Papavero N (2003) Uma breve história dos morcegos vampiros (Chiroptera, Phyllostomidae, Desmodontinae) no Brasil Colônia. Arquivos de Zoologia 43: 109–142.
- Thomas O (1899) List of the mammals obtained by Dr. G. Franco Grillo in the province of Parana, Brazil. Annali del Museo Civico di Storia Naturale di Genova 20: 546–549.
- Thomas O (1928) The Godman–Thomas expedition to Peru—VIII. On mammals obtained by Mr. Hendee at Pebas and Iquitos, Upper Amazons. Annals and Magazine of Natural History: Series 10 2: 285–294. <https://doi.org/10.1080/00222932808672879>
- Torquetti CG, Assis MAC, Carmo SSA, Talamoni SA (2013) Observations of a diurnal roost of the white-winged vampire bat *Diaemus youngi* in a karstic area of southeastern Brazil. Chiroptera Neotropical 19: 1147–1150.
- Trajano E (1987) Fauna cavernícola brasileira: composição e caracterização preliminar. Revista Brasileira de Zoologia 3: 533–561. <https://doi.org/10.1590/S0101-81751986000400004>
- Turner DC (1975) The vampire bat: a field study in behavior and ecology. Baltimore, John Hopkins University Press, 145 pp.
- Urbietta GL, Siqueira TYS, Gracioli G (2017) White-winged Vampire Bat, *Diaemus youngi* (Jentink, 1893) (Mammalia, Chiroptera): range extension in the Cerrado biome and new locality in Mato Grosso do Sul, southwestern Brazil. Check List 13: 2128. <http://doi.org/10.15560/11.5.1765>
- Uieda W, Chaves ME (2005) Bats From Botucatu Region, State of São Paulo, Southeastern Brazil. Chiroptera Neotropical 11: 224–226.
- Urbano-Vidales G, Sánchez-Herrera O, Téllez-Girón G, Medellín L. RA (1987) Additional records of Mexican mammals. The Southwestern Naturalist 32: 134–137.
- Valdez R, LaVal RK (1971) Records of bats from Honduras and Nicaragua. Journal of Mammalogy 52: 247–250. <http://doi.org/10.2307/1378465>
- Vargas-Mena JC, Alves-Pereira K, Barros MAS, Barbier E, Cordero-

-Schmidt E, Lima SMQ, Rodríguez-Herrera B, Venticinque EM (2018) The bats of Rio Grande do Norte state, northeastern Brazil. *Biota Neotropica* 18: e20170417. <http://doi.org/10.1590/1676-0611-bn-2017-0417>

Vieira CO da C (1942) Ensaio monográfico sobre os quirópteros do Brasil. *Arquivos de Zoologia* 3: 219–471.

Vizotto LD, Taddei VA (1973) Chave para determinação de quirópteros brasileiros. *Boletim de Ciências da Faculdade de Filosofia, Ciências e Letras* 1: 1–72.

Wenzel RL, Tipton VJ, Kiewlicz A (1966) The streblid batflies of Panama (Diptera Calypterae: Streblidae). *Fieldiana: Special Publications* 9: 405–675.

Wilkinson GS (1984) Reciprocal food sharing in the vampire bat. *Nature* 308: 181–184. <http://doi.org/10.1038/311525a0>

Zanon CMV, Reis NR (2007) Bats (Mammalia, Chiroptera) in the Ponta Grossa region, Campos Gerais, Paraná, Brazil. *Revista Brasileira de Zoologia* 24: 327–332. <https://doi.org/10.1590/S0101-81752007000200010>

Appendix

Table A1. Gazetteer of all reported occurrences of *Diaemus youngii* (Jentink, 1893).

| Country | State/province | Locality | Latitude | Longitude | Reference | Page | Coll. number |
|-----------|--------------------|--|-------------|--------------|--------------------------------|------|----------------|
| Argentina | Jujuy | Agua Salada | 23°49'00" S | 064°36'00" W | Barquez 1984 | 67 | FML 1343 |
| | Misiones | Bonpland | 27°29'00" S | 055°29'00" W | Kwon and Gardner 2008 | 222 | — |
| | Salta | Arroyo Arrazayal | 22°39'51" S | 064°25'31" W | Barquez et al. 2011 | 17 | CML 7700 |
| Bolivia | Beni | 15 km above Horquilla, Rio Machupo | 12°41'00" S | 064°32'00" W | Anderson et al. 1982 | 15 | — |
| | Beni | 4 km S San Joaquin, San Juan | 13°06'00" S | 064°49'00" W | Anderson et al. 1982 | 15 | — |
| | Beni | Estancia Yutiole | 13°15'00" S | 064°49'00" W | Anderson et al. 1982 | 16 | — |
| | Beni | Rio Itenez, 4 km above Costa Marquez | 12°29'00" S | 064°15'00" W | Anderson et al. 1982 | 14 | — |
| | La Paz | Pasto Grande | 16°36'00" S | 067°29'00" W | Kwon and Gardner 2008 | 222 | — |
| | Santa Cruz | Buena Vista | 17°27'00" S | 063°40'00" W | Kwon and Gardner 2008 | 222 | — |
| Brazil | Acre | Parque Nacional da Serra do Divisor | 07°27'30" S | 073°43'55" W | Nogueira et al. 1999 | 366 | — |
| | Alagoas | Porto Calvo | 09°04'00" S | 035°24'00" W | Kwon and Gardner 2008 | 222 | — |
| | Amapá | Santa Luzia do Pacuí | 00°30'00" N | 051°40'00" W | Peracchi et al. 1984 | 97 | ALP |
| | Amazonas | Biological Dynamics of Forest Fragments Project (BDFFP), 80 km N Manaus | 02°24'26" S | 059°43'40" W | Bernard 2001 | 119 | INPA |
| | Amazonas | Manaus | 03°06'00" S | 060°01'00" W | Mok and Lacey 1980 | 339 | — |
| | Amazonas | Parque Nacional do Jaú | 01°50'00" S | 062°50'00" W | Barnett et al. 2006 | 108 | — |
| | Amazonas | Rio Aripuanã | 06°08'13" S | 060°11'37" W | Bobrowiec 2012 | 280 | INPA |
| | Amazonas | Rio Madeira | 05°27'58" S | 060°45'07" W | Bobrowiec 2012 | 280 | INPA |
| | Bahia | Baía do Rio São Francisco (Médio) | 13°42'25" S | 042°51'00" W | Sá-Neto and Marinho-Filho 2013 | 91 | DCN/UESB |
| | Bahia | Mascote, Fazenda São José | 15°34'05" S | 039°17'07" W | Falcão 2007 | 330 | DE 606 |
| | Distrito Federal | Brazlândia, Gruta do Sal (DF-005), Fazenda Palestina | 15°30'00" S | 048°10'00" W | Aguiar et al. 2006 | 895 | — |
| | Distrito Federal | Sobradinho, Área de Proteção Ambiental Cafuringa | 15°35'00" S | 048°01'00" W | Aguiar and Antonini 2016 | 3 | — |
| | Goiás | Monte Alegre de Goiás | 13°15'23" S | 046°54'02" W | This Study | - | — |
| | Maranhão | Área de Proteção Ambiental Municipal do Inhamum | 04°53'30" S | 043°24'53" W | Olímpio et al. 2016 | 3 | — |
| | Mato Grosso | Cláudia, Renato River | 11°24'00" S | 055°02'00" W | Pedroso et al. 2018 | 155 | MZUSP 35713 |
| | Mato Grosso | Fazenda Acurizal | 17°45'00" S | 057°37'00" W | Schaller 1983 | 11 | — |
| | Mato Grosso do Sul | Aquidauana | 19°33'58" S | 055°30'43" W | Oliveira et al. 2012 | 36 | UNIDERP |
| | Mato Grosso do Sul | Caimã Forest and Resort | 19°18'13" S | 056°01'25" W | Urbieta et al. 2017 | 3 | — |
| | Mato Grosso do Sul | Campo Grande, Lagoa da Cruz, Instituto São Vicente | 20°23'08" S | 054°36'27" W | Urbieta et al. 2017 | 1 | ZUFMS CHI02310 |
| | Mato Grosso do Sul | Coxim, Serra Coxim | 18°35'21" S | 054°48'13" W | Urbieta et al. 2017 | 3 | — |
| | Mato Grosso do Sul | Miranda-Abobral | 19°46'55" S | 056°40'38" W | Oliveira et al. 2012 | 36 | UNIDERP |
| | Mato Grosso do Sul | Nhecolândia | 18°47'28" S | 056°02'03" W | Oliveira et al. 2012 | 36 | UNIDERP |
| | Mato Grosso do Sul | Nhecolândia, Fazenda Rio Negro | 19°34'22" S | 056°14'36" W | Silva and Graciolli 2013 | 177 | — |
| | Mato Grosso do Sul | Nhecolândia, Fazenda Rio Negro | 19°34'22" S | 056°14'36" W | Herrera et al. 2011 | 383 | — |
| | Mato Grosso do Sul | Paraguay river | 19°13'00" S | 057°29'00" W | Bordignon and Shapiro 2018 | 259 | — |
| | Minas Gerais | Esmeraldas | 19°45'57" S | 044°17'09" W | Urbieta et al. 2017 | 3 | — |
| | Minas Gerais | Lagoa Santa | 19°42'44" S | 043°56'32" W | Torquetti et al. 2013 | 1148 | MZ 270 |
| | Minas Gerais | Uberaba | 19°44'50" S | 047°56'21" W | Stutz et al. 2004 | 190 | — |
| | Minas Gerais | Uberlândia | 18°55'08" S | 048°16'39" W | Urbieta et al. 2017 | 3 | — |
| | Pará | Belém | 01°27'21" S | 048°29'15" W | Handley 1967 | 213 | — |
| | Pará | Belém, Área de Pesquisas Ecológicas do Guamá, e áreas do Instituto de Pesquisa e Experimentação Agropecuárias do Norte | 02°39'00" S | 055°22'00" W | Kalko and Handley 2001 | 322 | — |
| | Pará | Melgaço, Estação Científica Ferreira Penna | 01°48'00" S | 050°43'00" W | Marques-Aguiar et al. 2003 | 2 | — |
| | Pará | Rio Xingu, Usina Hidrelétrica Belo Monte | 03°20'00" S | 052°00'00" W | Marques-Aguiar et al. 2009 | 2 | — |
| | Pará | Santarém | 02°24'52" S | 054°42'36" W | Piccinini 1974 | 20 | — |
| | Paraíba | Araruna, Parque Estadual Pedra da Boca, Mata Seca | 06°27'43" S | 035°41'21" W | Feijó et al. 2010 | 723 | UFPB 5573 |
| | Paraíba | Reserva Biológica Guaribas | 06°48'18" S | 035°04'60" W | Feijó et al. 2016 | 65 | UFPB 7456 |
| | Paraná | Bacia do Rio Tibagi | 24°06'07" S | 050°53'06" W | Reis et al. 2002 | 255 | — |

Table A1. Continued.

| Country | State/province | Locality | Latitude | Longitude | Reference | Page | Coll. number |
|---------------|------------------------|--|-------------|--------------|-------------------------------|------|--------------|
| Brazil | Paraná | Cerro Azul | 24°49'00" S | 049°15'00" W | Graciolli and Carvalho 2001 | 935 | — |
| | Paraná | Londrina, Parque Estadual Mata dos Godoy | 23°27'00" S | 051°15'00" W | Reis et al. 2003 | 228 | — |
| | Paraná | Palmeira | 25°25'00" S | 050°00'00" W | Thomas 1899 | 547 | — |
| | Paraná | Parque Nacional do Iguaçu | 25°36'00" S | 054°35'00" W | Sekiama et al. 2001 | 752 | LZUEL |
| | Paraná | Ponta Grossa, Campos Gerais | 25°50'58" S | 050°09'30" W | Zanon and Reis 2007 | 329 | — |
| | Paraná | Roça Nova | 25°28'00" S | 048°58'00" W | Miller 1906 | 84 | USNM 140769 |
| | Pernambuco | Ipojuca, Usina Salgado, Mata do Mingú | 08°31'29" S | 035°03'26" W | Sotero-Caio et al. 2011 | 158 | — |
| | Pernambuco | Jaqueira, RPPN Frei Caneca | 08°42'37" S | 035°50'01" W | Silva et al. 2010 | 88 | — |
| | Pernambuco | São Lourenço da Mata, Estação Ecológica do Tapacurá | 08°02'00" S | 035°13'00" W | Mares et al. 1981 | 111 | — |
| | Piauí | Picos | 07°04'39" S | 041°28'02" W | Pinto and Bento 1986 | 32 | — |
| | Piauí | São Miguel do Tapuio | 05°43'13" S | 041°26'28" W | Castilho et al. 2010 | 1335 | — |
| | Rio de Janeiro | Angra dos Reis | 23°00'00" S | 044°18'00" W | Bolzan et al. 2010 | 589 | — |
| | Rio de Janeiro | Angra dos Reis, Ilha da Gipóia | 23°02'20" S | 044°21'35" W | Costa and Esbérard 2011 | 741 | — |
| | Rio de Janeiro | Baía de Sepetiba, Mangaratiba, Ilha de Marambaia, Praia Grande | 22°04'26" S | 043°57'56" W | Costa et al. 2008 | 218 | LDM 4314 |
| | Rio de Janeiro | Barra Mansa | 22°32'27" S | 044°10'38" W | Peracchi and Albuquerque 1971 | 410 | — |
| | Rio de Janeiro | Carmo | 21°55'00" S | 042°36'00" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | Casimiro de Abreu, Morro de São João | 22°29'96" S | 041°58'92" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | Guapiaçú, Reserva Ecológica de Guapiaçú | 22°25'53" S | 042°45'20" W | Souza et al. 2015 | 14 | MN 79877 |
| | Rio de Janeiro | Guapimirim, Parada Modelo | 22°33'24" S | 042°55'71" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | Mangaratiba | 22°56'34" S | 044°02'26" W | Bolzan et al. 2010 | 589 | — |
| | Rio de Janeiro | Mangaratiba, Fazenda Terras do Sahy | 22°56'00" S | 044°00'00" W | Costa and Esbérard 2011 | 741 | — |
| | Rio de Janeiro | Mangaratiba, Ilha de Marambaia | 23°04'03" S | 043°53'14" W | Costa and Esbérard 2011 | 741 | — |
| | Rio de Janeiro | Miracema | 21°24'55" S | 042°11'39" W | Barros et al. 2008 | 684 | — |
| | Rio de Janeiro | Quissamã, Parque Nacional Restinga de Jurubatiba | 22°06'14" S | 041°28'12" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | Rio de Janeiro, Barra da Tijuca | 23°00'22" S | 043°17'49" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | Rio de Janeiro, Barra de Guaratiba | 22°59'45" S | 043°32'21" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | RPPN Fazenda Bom Retiro | 22°27'13" S | 042°18'29" W | Menezes Junior et al. 2015 | 272 | — |
| | Rio de Janeiro | Seropédica, Campus da Universidade Federal Rural do Rio de Janeiro | 22°45'54" S | 043°41'12" W | Peracchi and Albuquerque 1986 | 67 | — |
| | Rio de Janeiro | Três Rios, Bemposta | 22°07'00" S | 043°12'20" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | Valença, Refúgio da Vida Silvestre da Serra da Concórdia | 22°22'18" S | 043°47'23" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | Volta Redonda | 22°30'27" S | 044°05'40" W | Costa et al. 2008 | 218 | — |
| | Rio de Janeiro | Volta Redonda, Parque Natural Municipal Fazenda Santa Cecília do Ingá | 22°27'34" S | 044°04'51" W | Pereira et al. 2013 | 1018 | — |
| | Rondônia | Porto Velho | 08°46'00" S | 063°54'00" W | Tavares et al. 2017 | 96 | — |
| | Rondônia | Porto Velho, Abunã | 09°35'00" S | 065°03'00" W | Pedroso et al. 2018 | 155 | MZUSP 35712 |
| | São Paulo | Bilac | 21°24'16" S | 050°28'30" W | Garbino 2016 | 82 | DZSJRP 10705 |
| | São Paulo | Botucatu | 22°59'20" S | 048°26'37" W | Uieda and Chaves 2005 | 225 | — |
| | São Paulo | Grota Mirassol | 20°46'00" S | 049°28'00" W | Garbino 2016 | 82 | DZSJRP 2556 |
| | São Paulo | Guararema | 23°25'00" S | 046°01'00" W | Garbino 2016 | 82 | DZSJRP 15156 |
| | São Paulo | Iporanga, Parque Estadual Turístico Alto do Ribeira, Gruta do Alambari de Baixo (SP-012) | 24°33'15" S | 048°39'55" W | Trajano 1987 | 538 | — |
| | São Paulo | Itajobi | 21°19'00" S | 049°03'00" W | Garbino 2016 | 82 | DZSJRP 15025 |
| | São Paulo | Santa Gertrudes, Fazenda Paraguassu | 22°27'00" S | 047°32'00" W | Sazima and Uieda 1980 | 102 | ZUEC 1001 |
| | São Paulo | São José do Rio Preto | 20°49'11" S | 049°22'45" W | Garbino 2016 | 82 | DZSJRP 16978 |
| | São Paulo | São Paulo | 23°32'52" S | 046°38'10" W | Vieira 1942 | 380 | MZUSP 4036 |
| | São Paulo | São Sebastião, Barra do Una | 23°46'00" S | 045°45'00" W | Garbino 2016 | 82 | MZUSP 9462 |
| | São Paulo | Uchoa | 20°57'00" S | 049°10'00" W | Garbino 2016 | 82 | DZSJRP 16615 |
| | Tocantins | Couto Magalhães, near electric power transmission line | 08°21'00" S | 049°10'00" W | Pedroso et al. 2018 | 156 | MZUSP 35358 |
| | Tocantins | Estação Ecológica Serra Geral do Tocantins, Jalapão | 10°33'00" S | 046°45'00" W | Gregorin et al. 2011 | 303 | EG 323 |
| Colombia | Putumayo | Río Mecaya | 00°28'00" N | 075°20'00" W | Wenzel et al. 1966 | 600 | — |
| | Valle del Cauca | Río Raposo | 03°38'00" N | 077°05'00" W | Wenzel et al. 1966 | 600 | — |
| Costa Rica | Guanacaste | Finca La Pacífica | 10°27'16" N | 085°07'41" W | Gardner et al. 1970 | 723 | LSUMZ 22533 |
| Ecuador | Esmeraldas | San Lorenzo | 01°17'00" N | 078°50'00" W | Pinto et al. 2007 | 244 | TTU 85358 |
| El Salvador | La Paz | 1 mi N La Herradura | 13°21'52" N | 088°57'00" W | Owen et al. 1990 | 418 | — |
| | La Paz | 3 mi NW La Herradura | 13°22'51" N | 088°58'53" W | Greenbaum and Jones 1978 | 5 | TTU 27591 |
| French Guiana | Cayenne | Paracou | 05°16'00" N | 052°55'00" W | Bernard and Fenton 2002 | 1137 | — |
| Guyana | East Berbice-Corentyne | Upper Canje Creek | 05°07'00" N | 057°37'00" W | Jentink 1893 | 282 | RNH 12088 |
| | Potaro-Siparuni | Iwokrama Forest | 04°40'00" N | 058°41'00" W | Bernard and Fenton 2002 | 1137 | — |
| | Rupunini | Kanunu Mountains | 03°22'00" N | 059°30'00" W | Barnett et al. 2006 | 125 | — |

Table A1. *Continued.*

| Country | State/province | Locality | Latitude | Longitude | Reference | Page | Coll. number |
|-------------------|---|--|-------------|--------------|----------------------------|------|------------------|
| Mexico | Chiapas | Ejido Benemérito de Las Américas | 16°31'02" N | 090°39'11" W | Urbano-Vidales et al. 1987 | 135 | — |
| | Mazu | Montes Azules | 16°50'00" N | 091°30'00" W | Bernard and Fenton 2002 | 1137 | — |
| Nicaragua | Managua | 0.75 mi N Masachapa | 11°47'24" N | 086°31'00" W | Greenbaum and Jones 1978 | 5 | TTU 30667 |
| | South Caribbean Coast Autonomous Region | 3 km NW Rama | 12°10'43" N | 084°13'12" W | Greenbaum and Jones 1978 | 5 | TTU 30668 |
| Panama | Bocas del Toro | Isla Bastimentos | 09°21'00" N | 082°12'00" W | Wenzel et al. 1966 | 600 | — |
| | Guna Yala | Río Armilla | 09°02'40" N | 078°01'14" W | Wenzel et al. 1966 | 496 | — |
| | Alto Paraguay | Estancia General Díaz, 100 km W Fuerte Olimpo | 21°08'00" S | 058°30'00" W | López-González et al. 1998 | 42 | FMNH 145266 |
| | Alto Paraguay | Estancia Parra Cue | 02°05'54" S | 047°53'31" W | López-González et al. 1998 | 42 | TTU 75276 |
| | Boquerón | Parque Nacional Defensores del Chaco, Cruce Cuatro de Mayo, destacamento Patricio Colmán | 20°13'00" S | 060°12'00" W | López-González et al. 1998 | 42 | MNHNP 823 |
| | Presidente Hayes | Estancia La Victoria | 23°29'02" S | 058°34'47" W | López-González et al. 1998 | 42 | TTU 75273 |
| | Presidente Hayes | Estancia Loma Pora | 23°29'56" S | 057°32'53" W | López-González et al. 1998 | 42 | TTU 75278 |
| Peru | Huánuco | Cueva de Castillo | 09°07'00" S | 075°58'00" W | Bowles et al. 1979 | 7 | — |
| | Huánuco | Cueva de Lechuzas | 09°20'00" S | 076°01'00" W | Bowles et al. 1979 | 7 | — |
| | Huánuco | Panguana Biological Station | 09°37'00" S | 074°56'00" W | Hutterer et al. 1995 | 5 | — |
| | Loreto | Colonia Calleria, 20 km from Río Ucayali | 08°06'00" S | 074°36'00" W | Wenzel et al. 1966 | 600 | — |
| | Loreto | Pebas | 03°20'00" S | 071°49'00" W | Thomas 1928 | 288 | NHM 1928.7.21.64 |
| | Tumbes | Distrito Pampas de Hospital, Angostura | 03°45'23" S | 080°23'15" W | Pacheco et al. 2007 | 412 | MUSM 22132 |
| | Ucayali | Yarinacocha | 08°15'00" S | 074°43'00" W | Sanborn 1949 | 282 | — |
| Trinidad & Tobago | Trinidad | Barataria | 10°39'00" N | 061°28'00" W | Kwon and Gardner 2008 | 222 | USNM 536938 |
| | Trinidad | La Brea | 10°15'00" N | 061°37'00" W | McBee et al. 1985 | 404 | TTU 26888 |
| | Trinidad | Las Cuevas | 10°46'00" N | 061°23'00" W | Greenhall 1963 | 397 | — |
| | Trinidad | Majuba Road, Petit Valley | 10°42'00" N | 061°32'00" W | Brennan 1967 | 154 | — |
| | Trinidad | Maracas Valley | 10°41'00" N | 061°24'00" W | McBee et al. 1985 | 404 | TTU 5232 |
| Venezuela | Amazonas | San Juan de Manapiare | 05°15'00" N | 066°05'00" W | Bernard & Fenton 2002 | 1137 | — |
| | Bolívar | Canaima National Park | 06°10'00" N | 062°30'00" W | Barnett et al. 2006 | 125 | — |
| | Delta Amacuro | Araguaimujo | 08°54'00" N | 061°32'00" W | Kwon and Gardner 2008 | 222 | — |
| | Falcón | 6 km SE Capatárida | 11°07'00" N | 070°35'00" W | Kwon and Gardner 2008 | 222 | — |
| | Nueva Esparta | El Valle | 10°59'00" N | 063°52'00" W | Kwon and Gardner 2008 | 222 | — |
| | Zulia | El Panorama, Río Aurare | 10°37'00" N | 071°25'00" W | Osgood 1912 | 63 | — |